

## Society for the Study of Amphibians and Reptiles

---

A New Species of Testudo (Testudines: Testudinidae) from the Middle East, with Implications for Conservation

Author(s): Jarmo Perälä

Source: *Journal of Herpetology*, Vol. 35, No. 4 (Dec., 2001), pp. 567-582

Published by: [Society for the Study of Amphibians and Reptiles](#)

Stable URL: <http://www.jstor.org/stable/1565894>

Accessed: 12/07/2011 16:17

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ssar>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



*Society for the Study of Amphibians and Reptiles* is collaborating with JSTOR to digitize, preserve and extend access to *Journal of Herpetology*.

thread snake, *Leptotyphlops scutifrons* (Serpentes, Leptotyphlopidae). J. Zool. Lond. 250:321–327.

———. 2000b. Life underground: food habits and reproductive biology of two amphisbaenian species (*Monopeltis anchietae* and *Zygaspis quadrifrons*) from southern Africa. J. Herpetol. 34:510–516.

YEADON, R. R. 1991. *Typhlops bibronii*, Bibron's blind snake: size and reproduction. J. Herpetol. Assoc. Afr. 39:23.

Accepted: 21 January 2001.

*Journal of Herpetology*, Vol. 35, No. 4, pp. 567–582, 2001  
Copyright 2001 Society for the Study of Amphibians and Reptiles

## A New Species of *Testudo* (Testudines: Testudinidae) from the Middle East, with Implications for Conservation

JARMO PERÄLÄ

School of Biological Sciences, University of Bristol, Woodland Road, Bristol BS8 1UG, UK;  
E-mail: j.perala@bristol.ac.uk

**ABSTRACT.**—Examination and analysis of over 140 specimens of the endangered tortoise *Testudo kleinmanni* from all areas within its historical distribution on mainland Africa and the Levant confirmed the existence of two allopatric species in this region. A new tortoise species, occurring east of the Nile delta, and restricted today to the Negev desert (Israel) and to northern Sinai (Egypt), is described. The new species has typically a rounded, considerably wide midbody and a very wide posterior carapace combined with very narrow vertebrals, a relatively modest elevation of the anterior plastron lobe, a gently sloping upper carapacial arch, which is posteriorly depressed, relatively flared, serrated edges of free marginals, a flared supracaudal in both sexes, usually not running in parallel with the posterior carapacial arch, and a short supracaudal in females, among other readily verifiable characteristics. Reflecting these relatively great differences in shell morphology, the new species differs from *T. kleinmanni* by 17 (male) and 18 (female) morphometric character ratios. The new species is additionally fully diagnosable by using Principal Component Analysis and Linear Discriminant Function Analysis with cross-validation. The previous lectotype designation for *T. kleinmanni* was not based on a syntype and is, therefore, invalid. A new lectotype is designated. Both species are endangered because of loss of habitat and other anthropogenic factors. The new species is additionally threatened by an introduction programme involving the release of confiscated *T. kleinmanni* into northern Sinai.

The Egyptian tortoise *Testudo kleinmanni* Lortet, 1883—the smallest representative of circum-mediterranean *Testudo*—was described from material collected by E. Kleinmann in 1875 near Alexandria in Egypt (Lortet, 1883, 1887). However, the name was considered a junior synonym of *Testudo leithii* Günther, 1869 until the publications by Mertens and Wermuth (1955) and Loveridge and Williams (1957). These authors concluded independently that the name *leithii* was already preoccupied by a fossil pelomedusid turtle described from India in 1852. Additionally, the type of *Testudo leithii* was claimed to originate from “Sindh, Baluchistan” (Günther, 1869), modern northwestern Pakistan, where the species has never been found again. Loveridge and Williams (1957) considered this type locality an error, but this geographic aspect already led Lortet (1887), who knew the description but had not seen the type, to doubt the

type locality of *leithii* and to speculate that *kleinmanni* and *leithii* might represent the same species. Lortet (1883) described *T. kleinmanni* without designating a holotype, but some of the syntypes still survive in a few collections, notably in the Lyon museum where he worked. Mertens (1967) listed, or designated (Iverson, 1992), a lectotype (SMF 7810; semiadult male; Alexandria; leg. and don. Lortet 1881) citing Boettger (1893), who already referred to the same specimen as “Typus von *T. kleinmanni*,” which could either be taken as an earlier lectotype designation according to Article 74 of the old ICZN Code (Anonymous, 1964) or as listing of a presumed syntype. In any case, the only real (physical and traceable) voucher specimens on which Lortet's (1883) description is based, that is, syntypes according to the Articles 72 and 73 of the old Code (Anonymous, 1964)—as indicated by two accounts (Lortet, 1883, 1887)—are those collected

by Kleinmann in the vicinity of Alexandria and donated to the author in 1875 (Lortet, 1887). Hence, SMF 7810 [specimen "3175 a" of Boettger (1893)] cannot represent a syntype as it was collected only in 1881 by Lortet himself, making any lectotype designation based on this specimen—whether by Boettger or Mertens—invalid, according to the old Code, Article 74(a)(i) (Anonymous, 1964). The type locality of *T. kleinmanni* is "dans les sables de la basse Égypte, surtout dans les environs d'Alexandrie" (Egypt), according to Iverson (1992), and is based on a direct quote from Lortet (1883).

Until recently, the range of *T. kleinmanni* was believed to extend along coastal regions from Tripolitania in western Libya (Iverson, 1992; Fritz and Buskirk, 1997; and confirmed by fieldwork by A. Pieh, pers. comm., June 2000), and Cyrenaica in eastern Libya in the west through mainland Egypt to north of El Arish on the Sinai Peninsula, Egypt, and northern Negev desert in Israel in the east (Lortet, 1883; Tristram, 1884; Calabresi, 1924; Flower, 1933; Loveridge and Williams, 1957; Werner, 1982, 1987, 1988; Buskirk, 1985; Geffen, 1990; Iverson, 1992; Baha El Din, 1994; Bringsøe and Buskirk, 1998; among others). Interestingly, the Nile delta seems to form a major distributional gap cutting the historical range of *T. kleinmanni* in half. There are no records from the delta between Alexandria and Damietta.

During fieldwork in the Negev desert and visits to collections in Jerusalem and Tel-Aviv, the suspicion arose that *T. kleinmanni* from east of the Nile delta might be specifically distinct from populations occurring in the west [in practice Libya, as populations on mainland Egypt are already extinct (Baha El Din, 1994)]. This study was initiated to test this hypothesis.

#### MATERIALS AND METHODS

*Summary of Material.*—Over 140 specimens of *T. kleinmanni* throughout the known range of the species were investigated. This sample is believed to contain almost all available material from Israel and the Sinai Peninsula. Museum specimens were examined in 10 scientific collections as listed in Appendix 1. Museum abbreviations are based on standards suggested by Leviton et al. (1985), with the exception of PCHP (Chelonian Research Institute, Oviedo; formerly: Peter C. H. Pritchard collection). Additionally, data were recorded from two living adults from the Negev desert, examined and released in the field by I. Siirilä, T. Koivu, and J. Perälä in 1997. The animals (field reference numbers JP 97.3.36-7) are two of three (2 adults, 1 juvenile) mentioned and depicted in Perälä (2000). Data from these individuals are deposited in BMNH.

*Grouping.*—The material was allocated into two geographical groups: (1) western, comprising animals from west of the Nile delta and (2) eastern, for specimens from east of the Nile delta. Specimens labeled "Egypt" without further detail of locality, were classified as western by morphological criteria after preliminary comparative examination of the material, except BMNH 1933.9.3.5-6, which were grouped into eastern category. Errors in allocation of specimens would subsequently show in multivariate analyses, and typically museum records discriminate between mainland Egypt (Egypt) and the Sinai Peninsula (Sinai).

The true origin of the type of *T. leithii* (BMNH 1947.3.4.35) was also of interest, and it was put initially into the western group on morphological grounds.

*Measurements.*—Quantitative data were recorded from 45 shell characters using calipers straight-line to the nearest 0.1mm. Additionally, with reference to one character (CU), a tape ruler was used to the nearest millimeter. Most parameters, as indicated with symbols, follow those published among others in Ernst and Lovich (1986‡), Iverson and McCord (1989+, 1997\*) and Ernst et al. (1998‡). All other measurements are defined below. Recorded characters included maximum (not midline) carapace length (CL; ‡, +, \*), maximum plastron length (PL; ‡, +, \*, †), midline (minimum) plastron length (PL-m), curved carapace length over the dome, from the anterior tip of nuchal scute to the posterior tip of supracaudal (CU), maximum midbody width within central bridge area not affected by a potential flaring of anterior or posterior marginals, that is, at marginals 5-6 (MI), maximum width of shell at posterior marginals 7-9 (MA), maximum (not midline) gular scute length (GU-l; +), maximum (combined, i.e., left plus right) gular scute width (GU-w; \*), gular scute height in a 90° angle to horizontal plastron level as the minimum distance between gular scute/humeral midline crossing and dorso-median gular lip surface (GU-h), maximum shell height (HE; +, \*), maximum inner width of anterior shell opening parallel to horizontal level (ASO-w), maximum inner height of anterior shell opening parallel to median axis (ASO-h), left minimum bridge length (BR; \*), maximum (combined) humeral scute width (and subsequent plastron scute widths) in a 90° angle to median axis (HUM-w), maximum (combined) pectoral scute width (PEC-w), maximum (combined) abdominal scute width (ABD-w), maximum (combined) femoral scute width (FEM-w), maximum (combined) anal scute width (AN-w), maximum nuchal scute length (NU-l; †), maximum nuchal scute width (NU-w; †), maximum gular midline (intergular) length (GU-m), left humeral mid-

line (interhumeral) seam length (HUM-m; +, \*), left pectoral midline (interpectoral) seam length (PEC-m; +, \*), left abdominal midline (interabdominal) seam length (ABD-m; +, \*), left femoral midline (interfemoral) seam length (FEM-m; +, \*), maximum anal midline (interanal) seam length (AN-m), maximum width of first vertebral scute (V1-w; †), maximum width of second vertebral scute (V2-w; †), maximum width of third vertebral scute (V3-w; †), maximum width of fourth vertebral scute (V4-w; †), maximum width of fifth vertebral scute (V5-w; †), maximum median length of first vertebral scute (V1-l; †), maximum median length of second vertebral scute (V2-l; †), maximum median length of third vertebral scute (V3-l; †), maximum median length of fourth vertebral scute (V4-l; †), maximum median length of fifth vertebral scute (V5-l; †), first costal length as the minimum straight-line distance between the anteriormost and posteriormost contact points with adjacent (normally first and fifth) marginal scutes (C1), costal two ventral length along marginals (C2), costal three ventral length along marginals (C3), costal four ventral length along marginals (C4), maximum dorsal width of supracaudal scute (SUP-d), maximum ventral (= max. total) width of supracaudal scute (SUP-v), maximum median length of supracaudal scute (SUP-l), maximum head width (HEAD), minimum distance between right eye and tympanum (EYE-TY), and, minimum distance between right eye and nostril (EYE-NO). Additionally, notes were made of body and individual scute forms, coloration, markings and potential abnormalities. All listed parameters could not always be recorded from every specimen examined due to shell injuries, abnormal growth/missing features, and human error (for further details, see below under "Data Analysis/Principal Components"). Some long-term wild-caught captives were used in the assessment of categorical character data (plastron markings) but excluded from morphometric analyses resulting from potential husbandry-related malformations, notably individuals from the Tel-Aviv University Expo Zoo—all from Israel according to Ora Kerman of TAU.

*Data Analysis.*—All statistical analyses were performed using MINITAB, Release 12.1 (Minitab, Inc.). Data, for every mensural character and for both populations separately, were preliminarily checked for normality (Anderson-Darling test) and for linearity by regression against maximum carapace length. Growth was allometric only in relatively small/young individuals [up to about 65 mm CL (males) and 75 mm CL (females), respectively], as expected and consistently isometric in subadult and adult specimens within both groups. Accordingly,

specimens with a maximum carapace length of less than those indicated above were excluded from further morphometric analyses (but were used in the assessment of categorical variation). Eventually, 99 specimens (36 ♂ + 40 ♀ western and 15 ♂ + 8 ♀ eastern) were compared for mensural character variation. The above material and additionally juveniles and originally (upon collection) crushed/damaged specimens (as listed in Appendix 1) were used in analyses of categorical data. According to inference from preliminary investigations, four individuals, MTKD D 34854 (western Libya), USNM 139093 [Gheminez, Cyrenaica (Libya)], TAU 12026 (Gevulot, Israel) and TAU 5635 (south of Gevulot, Israel) were omitted from further analyses because of their presumed hybrid status (intermediates between *T. kleinmanni* and tortoises currently classified into the *T. graeca* species complex). These putative hybrids differed from all other specimens by an atypical morphology or color pattern or both. Multivariate clustering of specimens from both populations was investigated using Principal Component Analysis (PCA). PCA components were calculated from the covariance matrix using data from 41 (males) and 40 (females) mensural characters. The remaining parameters—CU, PL-m, ASO-h (for females only), HEAD, EYE-TY, and EYE-NO—and four specimens (BMNH 1933.9.3.16; JP 97.3.37; MCZ 54045; PCHP 4447) were not taken into account owing to missing data. Sexes were analyzed separately. Discriminant Function Analysis (DFA), using linear discriminant function (Mahalanobis distance), and additionally cross-validation to compensate for an optimistic apparent error rate, was used to test how reliably specimens were classified into known groups using PCA scores from the first three principal components. Morphological character variation in 46 shell parameters among both groups was investigated using univariate analysis of variance (one-way ANOVA) with the significance level set at  $\alpha = 0.05$  and sexes treated separately. With respect to this character analysis, data were standardized for body size by division by maximum carapace length; thus the ratios eliminating the effect of size-related variation, and allowing for direct comparisons between individuals and populations. Despite some theoretical concerns associated with the use of ratios, this method of standardization has clearly been shown to produce unbiased and effective results in practice (Iverson and Graham, 1990) and is widely used in modern chelonian taxonomy (e.g., Cann and Legler, 1994; Iverson and McCord, 1994, 1997; Seidel et al., 1999). The normality of the residuals and equal variances were routinely checked relative to every character to be certain of the statistical validity of

the ANOVA results. The effect of geographical localities relative to the occurrence of distinct plastron markings or their combinations was checked using chi-square tests and the above significance level.

*Systematics.*—On theoretical grounds reviewed recently by Grismer (1999), in this study taxonomic inferences are made in accordance with the lineage-based evolutionary species concept (Simpson, 1961; Wiley, 1978; Frost and Hillis, 1990).

#### RESULTS AND DISCUSSION

In Principal Component Analysis, the first three components together account for 87.2% of the total variance (males) and 82.0% (females), with PC1 (interpreted as size) capturing by far the most information from the underlying data structure (males: eigenvalue 444.72, 81.7% of total variability; females: 420.0, 73.9%). Summary statistics and component loadings for PC1-3 are found in Table 1. PC scores for each population and gender are presented in Table 2. With respect to males, PC score plots (PC1+PC2 and PC2+PC3) show a clear separation of two completely nonoverlapping clusters corresponding to the geographical populations of *T. kleinmanni* from each side of the Nile delta (scores for PC1+PC2 are presented in Fig. 1). Variation on axis PC2 (interpreted to represent body form) is highly significant (ANOVA  $F_{1,49} = 129.39$ ,  $P < 0.0001$ ). As for females, plots of PC scores from PC1+PC2 (Fig. 2) also discriminate well between the two populations, and variation is again greatest on axis PC2 (ANOVA  $F_{1,44} = 69.59$ ,  $P < 0.0001$ ) although there is also statistically significant variation with respect to size (PC1; ANOVA  $F_{1,44} = 5.33$ ,  $P < 0.026$ ). Discriminant Function Analysis on scores from PC1-3 without or with cross-validation classified 100% of all individuals correctly into known groups irrespective of gender (Table 3) (the same result is achieved using raw character data). Additionally, as indicated by clustering of PC scores and DFA, there is strong evidence that the true geographic origin of the type of *Testudo leithii* Günther, 1869 lies within mainland Egypt (Fig. 2)—*T. kleinmanni* was not known from Libya until the early 20th century (Calabresi, 1924)—and that the type specimen was subsequently exported to Baluchistan or mixed up with material coming from there, as suggested by Lortet (1887).

Among males 17 and among females 18 significant to highly significant differences were found between the eastern and western population with respect to basic morphometric characters standardized for body length (Tables 4–6).

Pectoral blotches were present significantly

more often (80.9% of individuals) in animals from the eastern population than in specimens from the west (21.3%;  $\chi^2 = 37.759$ ,  $df = 1$ ,  $P < 0.0001$ ). A combination of abdominal and pectoral markings were present in 78.7% of animals from the east but only in 11.5% of individuals of western origin ( $\chi^2 = 49.725$ ,  $df = 1$ ,  $P < 0.0001$ ). Over one third (23 of 61, or 37.7%) of animals from the west showed no plastral markings whatsoever, as opposed to eastern tortoises in which some form of plastral pattern was present in every examined specimen ( $\chi^2 = 22.516$ ,  $df = 1$ ,  $P < 0.0001$ ).

The strong morphological heterogeneity of the two allopatric populations is interpreted as clear evidence for a divergent evolutionary history and the formation of two genealogical lineages within current *T. kleinmanni*. Though the ecology of *T. kleinmanni* has been researched in detail in Israel only, examination of relevant literature suggests the existence of discrepancies in ecological adaptations between the Israeli and mainland African populations (Baha El Din, 1994; Flower, 1933; Geffen and Mendelsohn, 1988, 1989, 1991; Schleich, 1989; Schleich et al., 1996). Because the eastern population is morphometrically distinct and geographically isolated from *T. kleinmanni* from mainland Egypt west of the Nile delta and Libya, it is here described as follows.

#### *Testudo wernerii* sp. nov.

Negev Tortoise

Figure 3

*Holotype.*—HJ 949, adult female, in alcohol. Collected by Haas, Werner et al., 12 March 1963, from sand dunes 14 km south of Be'er Sheva (northern Negev desert, Israel). This specimen was the first record of *T. kleinmanni* in Israel. [The species' occurrence as "*T. kleinmanni*" in the vicinity of Hebron and in Be'er Sheva in "Syria," or between Hebron and Be'er Sheva and in the Arabah south of the Dead Sea in "Palestine"—in a region that is nowadays Israel—was already known to both Lortet (1883) and Tristram (1884), respectively, but these observations were rebutted by Flower (1933) whose misinformed opinion was echoed by most subsequent authors (but see Bringsøe and Buskirk, 1998)].

*Paratypes.*—Seventeen specimens (locality data listed in Appendix 1): Males: BMNH 1900.2.8.11; BMNH 1933.9.3.12; HJ 16047; HJ 16048; TAU 5636; TAU 7468; TAU 8363; TAU 12022; TAU 12712; TAU 12769; TAU 14120; TAU 14126. Females: BMNH 1933.9.3; HJ 989; TAU 6295; TAU 11898; TAU 13397.

*Type Locality.*—Northern Negev desert, Israel.

*Distribution.*—*Testudo wernerii* spreads, historically, roughly from northern Negev near Be'er Sheva, and vicinities of Be'er Mash'abim, Holot

TABLE 1. Component loadings (eigenvectors) and summary statistics (eigenvalue, proportional and cumulative percentage of information captured) from eigenanalysis of the covariance matrix for first three Principal Components (PC1-3) from separate analyses of males and females of *Testudo kleinmanni* and *Testudo uermeri* sp. nov.

Variable	Males			Females			Variable	Males			Females		
	PC1	PC2	PC3	PC1	PC2	PC3		PC1	PC2	PC3	PC1	PC2	PC3
CL	-0.448	-0.063	-0.301	-0.420	-0.184	-0.111	AN-rn	-0.068	-0.130	0.126	-0.036	-0.006	0.057
PL	-0.360	-0.158	-0.509	-0.358	-0.106	0.004	PEC-w	-0.256	0.264	0.123	-0.285	0.339	-0.114
MI	-0.273	0.260	0.108	-0.299	0.205	-0.066	ABD-w	-0.258	0.221	0.111	-0.283	0.177	0.044
MA	-0.261	0.379	0.106	-0.292	0.352	0.057	V1-w	-0.063	-0.182	-0.018	-0.058	-0.086	-0.008
GUL-1	-0.017	-0.009	-0.110	-0.041	0.004	-0.068	V2-w	-0.084	-0.196	0.111	-0.088	-0.071	-0.017
GU-w	-0.046	-0.001	-0.114	-0.056	0.068	-0.071	V3-w	-0.114	-0.224	0.131	-0.109	-0.145	-0.014
GU-h	-0.030	-0.105	0.060	-0.029	-0.042	0.030	V4-w	-0.129	-0.205	0.119	-0.103	-0.265	0.003
HE	-0.192	-0.325	0.285	-0.184	-0.346	0.220	V5-w	-0.166	0.027	0.009	-0.116	-0.221	0.098
ASO-w	-0.162	0.171	0.038	-0.176	0.258	-0.117	V1-1	-0.073	-0.130	-0.083	-0.077	-0.068	-0.080
ASO-h	-0.057	0.075	0.051	*	*	*	V2-1	-0.082	-0.154	0.161	-0.082	-0.059	0.006
BR	-0.167	-0.141	-0.185	-0.185	-0.134	-0.054	V3-1	-0.100	-0.299	0.225	-0.088	-0.169	0.076
HUM-m	-0.140	0.108	0.045	-0.171	0.123	-0.029	V4-1	-0.089	-0.024	0.172	-0.093	-0.121	0.004
FEM-w	-0.174	-0.148	0.147	-0.186	0.071	0.188	V5-1	-0.113	0.084	0.064	-0.106	-0.124	-0.007
AN-w	-0.170	0.172	0.094	-0.144	0.038	0.133	C1	-0.141	-0.154	0.052	-0.136	-0.059	-0.029
NU-1	-0.021	-0.034	-0.080	-0.033	0.010	-0.036	C2	-0.082	0.008	-0.054	-0.999	0.037	-0.142
NU-w	-0.002	0.013	0.005	-0.007	0.022	-0.111	C3	-0.085	-0.054	0.077	-0.083	-0.063	-0.090
GU-m	-0.016	0.048	-0.075	-0.037	0.014	-0.082	C4	-0.103	-0.003	0.037	-0.081	-0.105	0.084
HUM-m	-0.055	-0.036	-0.015	-0.039	0.064	-0.035	SUP-d	-0.088	-0.093	0.181	-0.046	-0.090	0.048
PEC-m	-0.014	0.052	-0.253	-0.069	-0.146	-0.508	SUP-v	-0.142	-0.071	0.136	-0.120	-0.276	-0.123
ABD-m	-0.146	0.109	0.044	-0.108	0.071	0.684	SUP-1	-0.070	-0.125	-0.221	-0.040	-0.231	0.090
FEM-m	-0.032	-0.116	-0.225	-0.029	-0.057	-0.037	Eigenvalue	444.72	17.82	11.86	420.00	27.36	18.78
							Proportion	0.817	0.033	0.022	0.739	0.048	0.033
							Cumulative	0.817	0.850	0.872	0.739	0.787	0.820






FIG. 1. Principal Component scores from PC1 and PC2 for males of *Testudo kleinmanni* (plusses) and *Testudo wernerii* sp. nov. (open circles).




FIG. 2. Principal Component scores from PC1 and PC2 for females of *Testudo kleinmanni* (plusses) and *Testudo wernerii* sp. nov. (open circles). Type specimens are indicated as follows: HUI 949 (14 km south of Be'er Sheva, Israel), type of *T. wernerii* sp. nov. (solid circle); MG 42000414 (Alexandria, Egypt), designated lectotype of *T. kleinmanni* (solid square); BMNH 1947.3.4.35 ("Sindh, Baluchistan"), type of *Testudo leithii* (open square).

Haluza, Holot Agur, and areas SE of Dimona, further southwest along Mediterranean coastal regions on the Sinai Peninsula to the Nile delta in mainland Egypt. Today, it is almost entirely restricted to Israel. Lortet's (1887) reference to Damietta (modern Dumyāt), where he saw live animals (referred to as *T. kleinmanni*), cannot be substantiated by existing museum material, but is credible. A specimen (BMNH 1900.2.8.11) collected 1900 by Stanley Flower in Kantarah-il-Khastreh (possibly modern Al Qantarrah) at the Suez Canal represents the westernmost verifiable record for *T. wernerii*. Baha El Din (1992) reported a single individual from the southern shore of Lake Bardawil, and unconfirmed sightings of single animals from near El Gorah and 20 km southwest of El Arish, on northern Sinai. A single population was recently found on northern Sinai near the Israeli border (S. M. Baha El Din, pers. comm., 2000; J. R. Buskirk, pers. comm., 2000). The distributions of *T. kleinmanni* and *T. wernerii* correspond to great extent with those of the lacertid lizards *Acanthodactylus pardalis* (Lichtenstein, 1823) (coastal areas west of the Nile in Egypt to Cyrenaica, Libya) and the recently described *A. beershebensis* Moravec et al., 1999, a Negev endemic (Moravec et al.,

1999). Previously, *A. pardalis* was considered to occupy the pooled distribution of both species.

**Etymology.**—The new species is named in honor of Y. L. Werner of the Hebrew University of Jerusalem, one of the most active pioneers and true greats of Israeli and Middle Eastern herpetology, who has been over the years largely responsible for assembling the finest collection of herpetological specimens in the Middle East (HUI: Zoological Museum, Herpetology).

**Diagnosis.**—Although *T. wernerii* is superficially similar to *T. kleinmanni*, in practice it is readily distinguished from the latter especially (in dorsal and ventral view) by the rounded, considerably wide midbody combined curiously with narrow vertebrals and a wide posterior carapace, a relatively modest elevation of the anterior plastron lobe, a gently sloping upper carapacial arch which is relatively depressed posteriorly (in lateral view), the flared, dentate (serrated) free marginals, a flared supracaudal in both sexes not running in parallel with the pos-

TABLE 3. Summary of classification from Discriminant Function Analysis with cross-validation. Linear method for response: Species (sexes treated separately). Predictors: Principal Components 1–3 (PC1–3).

Classification group	Males		Classification group	Females	
	"True" group			"True" group	
	<i>kleinmanni</i>	<i>wernerii</i>		<i>kleinmanni</i>	<i>wernerii</i>
<i>kleinmanni</i>	35	0	<i>kleinmanni</i>	38	0
<i>wernerii</i>	1	15	<i>wernerii</i>	0	7
Total N	35	15	Total N	38	7
N correct	35	15	N correct	38	7
Proportion	1.000	1.000	Proportion	1.000	1.000



TABLE 4. Results of univariate analysis of variance of ratios of 17 (♂ ♂) and 18 (♀ ♀) characters to carapace length between males and females of *Testudo kleinmanni* (K) and *Testudo uermeri* sp. nov. (W). Sample sizes: Males: N = 36 (K) and N = 15 (W); Females: N = 40 (K) and N = 8 (W), except where indicated with an asterisk (\*).

Character	Males			Females			
	F	P	kleinmanni v uermeri	Character	F	P	kleinmanni v uermeri
CU	21.72	<0.0001	K > W *(K, N = 24; W, N = 4)	CL	4.88	0.032	K < W
MI	4.69	0.035	K < W	CU	22.03	<0.0001	K > W *(K, N = 28; W, N = 3)
MA	15.25	<0.0001	K < W	MI	11.39	0.002	K < W
GU-h	11.4	0.001	K > W	MA	18.71	<0.0001	K < W
HE	13.84	0.001	K > W	Gu-h	4.95	0.031	K > W
PEC-w	5.78	0.020	K < W	HE	12.61	0.001	K > W
ABD-w	5.36	0.025	K < W	ASO-w	7.85	0.007	K < W *(K, N = 40; W, N = 7)
FEM-w	7.31	0.009	K < W	HUM-w	5.49	0.024	K < W
AN-1	7.71	0.008	K > W	PEC-w	14.58	<0.0001	K < W *(K, N = 40; W, N = 7)
V1-w	19.96	<0.0001	K > W	ABD-w	10.8	0.002	K < W *(K, N = 40; W, N = 7)
V2-w	10.3	0.002	K > W	AN-w	4.07	0.049	K < W
V3-w	14.25	<0.0001	K > W	V3-w	4.83	0.033	K > W
V4-w	10.35	0.002	K > W	V4-w	13.49	0.001	K > W
V1-1	22.41	<0.0001	K > W	V5-w	9.06	0.004	K > W
V2-1	11.74	0.001	K > W	V3-1	9.51	0.003	K > W
V3-1	16.73	<0.0001	K > W	SUP-d	4.08	0.049	K > W
CI-1	6.04	0.018	K > W	SUP-1	19.69	<0.0001	K > W *(K, N = 24; W, N = 3)
				EYE-TYMP	9.32	0.005	K < W



TABLE 6. Descriptive statistics for 45 basic character ratios (and body length = CL in mm) for females of *Testudo kleinmanni* and *Testudo wernerii* sp. nov. Characters were standardized for CL. Abbreviations: Char = character; Mean = mean ratio; SE = standard error; SD = standard deviation; N = sample size.

Taxon	Char	Mean	SE	SD	N	Char	Mean	SE	SD	N	Char	Mean	SE	SD	N
<i>kleinmanni</i>	CL	111.495	1.363	8.620	40	NU-1	0.072	0.001	0.009	39	V2-1	0.221	0.002	0.015	40
<i>wernerii</i>		119.250	3.968	11.223	8	NU-w	0.075	0.003	0.007	8		0.212	0.004	0.012	8
<i>kleinmanni</i>	PL	0.875	0.003	0.022	40	GU-m	0.048	0.002	0.013	40	V3-1	0.214	0.002	0.014	40
<i>wernerii</i>		0.874	0.010	0.029	8	HUM-m	0.055	0.003	0.008	8		0.196	0.007	0.020	8
<i>kleinmanni</i>	PL-m	0.813	0.004	0.023	40	PEC-m	0.099	0.002	0.010	40	V4-1	0.205	0.002	0.013	40
<i>wernerii</i>		0.825	0.012	0.032	7	ABD-m	0.103	0.003	0.009	8		0.200	0.005	0.014	8
<i>kleinmanni</i>	CU	1.284	0.005	0.026	28	FEM-m	0.109	0.002	0.012	40	V5-1	0.198	0.003	0.019	4
<i>wernerii</i>		1.212	0.008	0.015	3	AN-m	0.116	0.010	0.028	8		0.194	0.003	0.009	8
<i>kleinmanni</i>	MI	0.718	0.003	0.020	40	PEC-w	0.119	0.008	0.023	40	C1	0.319	0.003	0.016	40
<i>wernerii</i>		0.747	0.011	0.031	8	FEM-w	0.277	0.006	0.018	8	C2	0.315	0.007	0.019	8
<i>kleinmanni</i>	MA	0.714	0.004	0.025	40	AN-w	0.155	0.003	0.017	40	C3	0.225	0.004	0.025	40
<i>wernerii</i>		0.755	0.009	0.024	8	PEC-m	0.277	0.006	0.018	8		0.230	0.003	0.010	8
<i>kleinmanni</i>	GU-1	0.110	0.001	0.009	40	ABD-w	0.074	0.003	0.017	40	C4	0.199	0.003	0.017	40
<i>wernerii</i>		0.113	0.003	0.008	8	V1-w	0.066	0.005	0.015	8	SUP-1	0.192	0.003	0.009	8
<i>kleinmanni</i>	GU-w	0.145	0.002	0.011	40	V2-w	0.155	0.003	0.017	40		0.183	0.002	0.014	40
<i>wernerii</i>		0.153	0.002	0.007	8	V3-w	0.143	0.013	0.038	8	SUP-d	0.176	0.005	0.013	8
<i>kleinmanni</i>	GU-h	0.105	0.001	0.008	39	V4-w	0.628	0.004	0.026	40		0.149	0.003	0.018	40
<i>wernerii</i>		0.09	0.003	0.009	8	V5-w	0.670	0.012	0.033	7		0.135	0.006	0.016	8
<i>kleinmanni</i>	HE	0.536	0.004	0.026	40	V1-1	0.654	0.003	0.020	40		0.255	0.004	0.027	40
<i>wernerii</i>		0.498	0.012	0.034	8	V2-1	0.683	0.012	0.030	7		0.236	0.013	0.035	8
<i>kleinmanni</i>	ASO-w	0.441	0.003	0.022	40	V3-1	0.209	0.002	0.014	40		0.160	0.003	0.016	40
<i>wernerii</i>		0.465	0.007	0.017	7	V4-1	0.199	0.006	0.017	8		0.133	0.003	0.010	8
<i>kleinmanni</i>	ASO-h	0.148	0.002	0.010	39	V5-1	0.250	0.002	0.013	40	HEAD	0.143	0.001	0.008	31
<i>wernerii</i>		0.150	0.004	0.013	8		0.245	0.005	0.015	8		0.142	0.002	0.005	5
<i>kleinmanni</i>	BR	0.476	0.002	0.013	40		0.290	0.002	0.016	40	EYE-TY	0.052	0.001	0.006	24
<i>wernerii</i>		0.471	0.008	0.022	8		0.277	0.005	0.013	8		0.063	0.005	0.005	3
<i>kleinmanni</i>	HUM-w	0.427	0.002	0.013	39		0.257	0.003	0.017	40	EYE-NO	0.044	0.001	0.005	26
<i>wernerii</i>		0.440	0.007	0.019	8		0.234	0.005	0.013	8		0.041	0.002	0.003	4
<i>kleinmanni</i>	FEM-w	0.456	0.003	0.018	40		0.267	0.003	0.017	40					
<i>wernerii</i>		0.467	0.006	0.018	8		0.248	0.003	0.007	8					
<i>kleinmanni</i>	AN-w	0.332	0.002	0.015	40		0.181	0.002	0.013	40					
<i>wernerii</i>		0.345	0.008	0.023	8				0.012	8					




FIG. 3. Holotype of *Testudo wernerii*, HUJ 949, adult female, 131.0 mm CL. Photograph by Aibi Niv (HUJ).

terior carapacial arch, and the short supracaudal of females.

*Description.*—A small-sized species of *Testudo*, carapace length up to 131.0 mm in females (mean 119.25 mm) and 106.4 mm in males (mean = 97.53 mm). Xiphiplastral kinesis is well developed in both sexes, although this feature is evidently not as pronounced as in *T. kleinmanni* in which latter species the posterior plastron lobe is exceptionally well hinged allowing for maximal flexibility. The interpectoral seam is longer than the interfemoral, as is also typical of *T. kleinmanni*. The anterior carapace rim between the first marginals is only modestly indented towards the nuchal scute, the carapace is usually relatively low domed (usually less than 50% of length in males) and it is highest at more or less between the anterior edge and the midpoint of the third vertebral scute. *T. kleinmanni* is typically highest at the posterior—or middle and posterior—portion of vertebral three. The upper part of the carapacial arch of *T. wernerii* is modestly developed, gently sloping and angulate or edged rather than rounded in anterior, posterior and lateral view, and the posterior part of the carapace is relatively depressed as opposed to a more rounded posterior part in *T. kleinmanni*. *Testudo wernerii* differs further from all other *Testudo* species by the following combination of characters (subject to individual variation): a pronounced flaring, reversion, dentation, and rounding of the free edges of all but the bridge marginals (features present

in a minor portion of Egyptian and Libyan animals only); a relatively great flaring to the rear of the supracaudal, which scute forms usually an angle with the fifth vertebral (in *T. kleinmanni* the supracaudal runs usually continuously parallel with V5); a marked reversion of the posterior edge of the supracaudal; a relatively narrow contact between fourth and fifth vertebrales; a relatively wide midbody; a relatively large maximum carapace width (at posterior marginals); a relatively low gular/epiplastral region combined with a well (but moderately in comparison with *T. kleinmanni*) elevated anterior plastron lobe in both sexes; a relatively wide anterior shell opening in females; a relatively wide plastron across humeral scutes in females; a relatively wide plastron across pectoral and abdominal scutes in both sexes; a relatively narrow plastron at femorals in males leaving the hind feet a comparably large space between the posterior plastron lobe and the adjacent marginals; a relatively broad combined anal scute width in females; a relatively short interanal seam in males; very narrow first through fourth vertebrales in males and relatively narrow third, fourth and fifth vertebrales in females [vertebral 1 has typically slightly rounded lateral sides only but not a "lyre shape" (Gmira, 1995) as common in *T. kleinmanni*]; relatively short first and second vertebrales in males; relatively short third vertebrales in both sexes; the tendency for rounding of the dorsal surface of the fifth vertebral scute; a relatively short first costal scute

in males; dorsally a relatively narrow, and medially a relatively short supracaudal scute in females and, a relatively long temporal region in females.

The extremities are relatively slender, featuring anteriorly five and posteriorly four thin and cylindrical, more or less transparent, claws. The front feet have bulky and subconical, mostly overlapping enlarged scales totalling three (proximal) to between four and six (outermost) in a longitudinal row (from elbow to claws) and three in a transverse row. These scales, of uniformly bright yellow to beige color, are strongly elevated anteriorly from the surface of the extremities (possibly as an adaptation to enable effective movement in loose sand). The apex of these scales tends to be generally more pointed than in *T. kleinmanni*. The throughout yellow to beige hind feet feature 2–3 swordlike enlarged scales at the heels, as is common in various species of *Testudo*. The tail is relatively long and slender, longer in males than in females, and without a horny tubercle at the tip. The thighs lack enlarged conical tubercles (spurs). The skin of soft parts is very thin and flexible, almost parchment-like and transparent.

Coloration of carapace in life is bright or dull yellow, typically with well-defined markings of dark brown to dark brownish-violet, very thin to medium broad bands (covering up to half of a given scute) running along the anterior and lateral margins of the vertebral scutes, and the anterior and lateral margins of the costals, much as in some but not most *T. kleinmanni*. Anterior edges of the marginal scutes feature often, but not necessarily, a dark and horizontally broad triangular pattern with the vertical apex located dorsally and the horizontal apex oriented posteriorly (the bridge marginals may be without any pattern). Such broad triangular markings on the marginals are rarely present in *T. kleinmanni*, in which species they are usually replaced by thin vertical bands or no bands at all. The basic coloration of the plastron is mostly bright or dull yellow. Dark brown, in most cases very well-defined triangular markings are typically present (97.8% of examined specimens) on the abdominal scutes, with the base at the anterior border and the apex oriented posteriorly. The anterior borders of the pectoral scutes feature usually (80.9% of examined specimens) a broader than long, dark brown fleck or semitriangular or rounded marking with the apex situated posteriorly. These markings are rarely present in one of the paired pectorals only. Additionally, according to the material examined, *T. werneri* seems to typically feature some kind of plastral pattern (100% of individuals examined); therefore, animals showing no plastral markings can be classified as *T. kleinmanni* with reasonable

confidence by coloration only. The basic coloration of soft parts, head and posterior leg surfaces in life vary from a very pale and bright yellow to a greenish or grayish yellow. The eyes are often bordered by darker scales, and an irregular spot of dark scales may be found on top of the head.

*Description of Holotype.*—A large adult female with relatively elevated vertebral and costal shields, with the posterior carapacial arch declining modestly toward the supracaudal, the scute of which is ventally broad, strongly reverted, and flared to the rear. Highest at one-third the length of vertebral three. Midbody relatively wide, widest at sixth marginals. Maximum body width at eighth marginals. Vertebrales and supracaudal/V5 contact relatively narrow. Lower posterior edges of all free marginals well serrated, reverted, and showing a relatively large degree of flaring. Bridge marginals feature a well-developed lateral keel. Mechanical injury visible (minor part broken off) on posterior free edge of left second marginal. Anterior plastron lobe greatly elevated (but very modestly in comparison with *T. kleinmanni*). Lateral sides of femoral scutes relatively straight from base to around three-fourths the length. Gular region well exposed from anterior plastron surface in ventral view. Anterior extremities with three longitudinal rows of very large, subconical and partially overlapping scales, totalling six on the outer surface, and 3–4 closer to the body. Basic coloration of shell in alcohol is greyish yellow, with dark brown markings. The vertebrales feature relatively broad dark bands covering the anterior and lateral borders of each scute, with dark pigment filling up to 50% of the scute surface (notably V1-3). Anterior and lower lateral edges of costals feature thin to medium broad markings, which in places become striated toward the center of each scute. The anterior borders of anterior and posterior marginals are pigmented with more or less well-defined triangular or semitriangular markings. The bridge marginals feature only shades of such markings. The lateral borders and anterior (except for immediate central) surface of the supracaudal also feature a broad dark band. The nuchal is unmarked. Well-defined dark brown and triangular markings are present on both pectoral and abdominal scutes, with the bases located at the extreme anterior borders of the respective scute. Soft parts and head are basically greyish to bright yellow. Aggregates of brownish scales are present around the eyes, and on top of the head.

*Dimensions of Holotype.*—Maximum carapace length 131.0 (all measurements in millimeters), maximum plastron length 119.5 (91.2% of CL), median (midline) plastron length 111.0 (84.7%

of CL), midline width within bridge area 96.5 (73.7% of CL), maximum width at posterior marginals, i.e. maximum total width 97.4 (74.4% of CL), maximum gular scute length 15.9 (12.1% of CL), combined gular scute width 20.1 (15.3% of CL), gular height 13.1 (10.0% of CL), maximum height 70.5 (53.8% of CL), inner width of anterior shell opening 63.3 (48.3% of CL), inner height of anterior shell opening 23.3 (15.5% of CL), length of bridge 61.1 (46.6% of CL), combined humeral scute width 60.7 (46.3% of CL), combined femoral scute width 60.7 (46.3% of CL), combined anal scute width 46.5 (35.5% of CL), nuchal scute length 11.1 (8.5% of CL), maximum (posterior) nuchal scute width 7.0 (5.3% of CL), gular midline length 14.7 (11.2% of CL), humeral midline length 14.9 (11.4% of CL), pectoral midline length 18.2 (13.9% of CL), abdominal midline length 35.6 (27.2% of CL), femoral midline length 10.1 (7.7% of CL), anal midline width 21.4 (16.3% of CL), combined pectoral scute width 85.4 (65.2% of CL), combined abdominal scute width 87.1 (66.5% of CL), first vertebral width 25.2 (19.2% of CL), second vertebral width 33.3 (25.4% of CL), third vertebral width 37.2 (28.4% of CL), fourth vertebral width 29.9 (22.8% of CL), fifth vertebral width 31.3 (23.9% of CL), first vertebral length 26.0 (19.8% of CL), second vertebral length 28.5 (21.8% of CL), third vertebral length 27.7 (21.1% of CL), fourth vertebral length 24.6 (18.8% of CL), fifth vertebral length 25.1 (19.2% of CL), first costal length 40.2 (30.7% of CL), second costal length 30.1 (23.0% of CL), third costal length 24.8 (18.9% of CL), fourth costal length 22.3 (17% of CL), dorsal supracaudal width 18.2 (13.9% of CL), ventral supracaudal width 36.1 (27.6% of CL), median supracaudal length 16.7 (12.7% of CL), maximum head width 18.6 (14.2% of CL), minimum distance between eye and tympanum 7.6 (5.8% of CL), and minimum distance between eye and nostril 5.9 (4.5% of CL).

*Mertens'/Boettger's Lectotype*.—The lectotype designations by Mertens (1967), or, depending on interpretation, Boettger (1893), are not valid, as discussed previously in this article; thus *T. kleinmanni* lacks a type specimen. Therefore, MG 42000414, an alcohol-preserved adult female and syntype from Alexandria (Egypt) in the original Lortet collection in Lyon, is hereby designated lectotype of *T. kleinmanni*. This specimen is presumably one of the individuals collected by Kleinmann in 1875, because it died a year later at the Lyon museum. Hence, all former syntypes of *T. kleinmanni* are considered paralectotypes of the same taxon.

*Description of Lectotype*.—A large adult female with a very slender (narrow) body and well-domed (rounded) upper carapacial arch and abruptly descending sides. Carapace is highest

at posterior region of third vertebral. Supracaudal is ventrally narrow, showing no great flaring to the rear, and runs almost parallel to fifth vertebral in lateral view. Supracaudal/V5 contact is relatively wide. Vertebrae are relatively broad, but V1 is very narrow anteriorly. Free marginals are only slightly flared and very modestly reverted, if at all, and not serrated except for M1,2 and 3. Anterior plastron lobe is well elevated from horizontal level. Basic shell color (in alcohol) is wooden yellow. Anterior and lateral borders of supracaudal, nuchal, all vertebrae and marginals, and anterior and lower lateral edges of the costals feature a relatively thin to medium broad band of brown pigmentation, which becomes striated toward the areolae. Only the third marginals show a hue of triangular pattern. The abdominals feature relatively poorly defined (delimited) triangular markings. The remaining plastral scutes are without any distinct pattern. Anterior extremities feature three longitudinal rows of very large, subconical and partially overlapping scales, totalling five on the outer surface and four closer to the body. Soft parts and head are grayish yellow, the head also with light brown, poorly delimited scutes on the top and snout, and dorsal, dark brown flecks in front of and behind the eyes, and in the posteriormost portion of the temple.

*Dimensions of Lectotype*.—CL: 114.8 (all measurements in millimeters), PL: 100.7 (87.7% of CL), PL-m: 93.7 (81.6%), MI: 79.5 (69.3%), MA: 80.5 (70.1%), GU-l: 13.1 (11.4%), GU-w: 18.1 (15.8%), GU-h: 11.5 (10.0%), HE: 59.0 (51.4%), ASO-w: 51.6 (44.9%), ASO-h: 15.7 (13.7%), BR: 53.4 (46.5%), HUM-w: 49.3 (42.9%), FEM-w: 49.7 (43.3%), AN-w: 36.7 (32.0%), NU-l: 8.8 (7.7%), NU-w: 6.4 (5.6%), GU-m: 12.7 (11.1%), HUM-m: 13.4 (11.7%), PEC-m: 17.9 (5.6%), ABD-m: 29.7 (25.9%), FEM-m: 5.6 (4.9%), AN-m: 17.3 (15.1%), PEC-w: 72.6 (63.2%), ABD-w: 72.2 (62.9%), V1-w: 24.4 (21.3%), V2-w: 28.6 (24.9%), V3-w: 32.6 (28.4%), V4-w: 27.0 (23.5%), V5-w: 28.4 (24.7%), V1-l: 20.5 (17.9%), V2-l: 25.8 (22.5%), V3-l: 23.9 (20.8%), V4-l: 22.2 (19.3%), V5-l: 23.2 (20.2%), C1: 38.8 (33.8%), C2: 28.8 (25.1%), C3: 24.6 (21.4%), C4: 19.6 (17.1%), SUP-d: 19.8 (17.2%), SUP-v: 29.1 (25.3%), SUP-l: 15.3 (13.3%), HEAD: 17.5 (15.2%), EYE-TY: 7.4 (6.4%), EYE-NO: 5.8 (5.1%).

*Conservation*.—The elevation of *T. kleinmanni* onto CITES I Appendix in 1995 essentially banned all legal trade in this species. However, this did not stop the illegal trade in animals within Egypt, despite a confiscation of more than 200 traded animals in the Cairo bazaars in 1997 (Highfield, 1997, and pers. comm.; Perälä, 2000). Some of these originally Libyan animals, that is, *T. kleinmanni*, have been released into an open-air enclosure and others introduced on an

island, at Lake Bardawil on northern Sinai as part of a conservation program run in cooperation by Egyptian wildlife authorities and various international nongovernmental organisations, and the Royal Netherlands Embassy in Cairo (Baha El Din, 1999; E. Wenman, pers. comm., July 2000). This ongoing reintroduction program—which was established despite the fact that *T. kleinmanni* from Libya and north-western Egypt, as opposed to those from northern Sinai and Israel, were known to have different ecological adaptations as cited above and discussed in Perälä (2000)—poses a major and immediate threat to the survival of *T. wernerii*. Even if individuals were to be released in fenced areas or on an island, this would not be a secure solution. Introduced animals held in “closed” environments or farms have the tendency to escape into the wild. The introduction of *T. kleinmanni* within the range of *T. wernerii* could contribute to the mixing (hybridization) of the two populations with devastating consequences and loss of biodiversity. However, hybridization is not the only serious threat posed by the introductions. Alien micro-organisms could also establish in new habitats. In similar circumstances, pathogens from released animals are suggested to have caused a large scale epidemic among wild *Gopherus agassizii* in Nevada and California (Jacobson et al., 1995).

In less than a hundred years, as predicted by Flower (1933), *T. kleinmanni* has become so rare that it is on the verge of extinction in the wild. The species is currently “technically” extinct in its former main distribution range in Egypt (Baha El Din, 1994) because of habitat destruction and fragmentation and other mainly anthropogenic factors discussed extensively in the same paper. *Testudo kleinmanni* is currently considered one of the most endangered chelonian species in the world as indicated by its CITES listing and IUCN Red List status (endangered). The discovery that *T. kleinmanni* is actually not a single taxon with a highly fragmented population structure but two separate species facing extinction should be taken into consideration in legislation and conservation efforts at national and international level. The protection of habitats occupied by *T. wernerii* on Sinai and in Israel is essential for its survival.

*Acknowledgments.*—This research was kindly supported by the Chelonian Research Foundation's (Lunenburg, MA) Linnaeus Fund Award, the London Natural History Museum Bioresource LSF Programme, and the Paris Muséum National d'Histoire Naturelle Parsyst Programme. Additional sponsorship for fieldwork was received from the Herpetological Society of Finland. I thank R. A. Avery and P. J. Miller

(School of Biological Sciences, University of Bristol) for their critical review of an earlier draft of the manuscript and their continuous support. I thank H.-H. Schleich and an anonymous reviewer, and also A. Bauer, for their critical comments and suggestions during the review process. Y. L. Werner's comments on part of the manuscript were much appreciated. I am additionally greatly indebted to the following persons for their help and support or for information: B. Andreas (MTKD), E. N. Arnold (BMNH), G. Babocsay (HUJ), S. Baha El Din (Cairo), B. Bordenave (MNHN), R. Bour (MNHN), H. Bringsøe (NHF, Denmark), J. Buskirk (Oakland), J. Camhi (HUJ), B. Clarke (BMNH), J. Clary (MG), T. Dayan (TAU), U. Fritz (MTKD), S. Hervet (MNHN), A. C. Highfield (Tortoise Trust, UK), O. Kerman (TAU), T. Koivu (University of Turku and Herpetological Society of Finland), France de Lapparent de Broin (MNHN), N. Lee (Bristol), C. McCarthy (BMNH), A. Niv (HUJ), E.-J. Obst (MTKD), A. Ohler (MNHN), O. Pauwels (MNHN), L. and S. Perälä (Helsinki), A. Pieh (MTKD and Leipzig University), P. C. H. Pritchard and family (PCHP), J. Rosado (MCZ), H. Seligmann (HUJ), I. Abdul Siirilä (Helsinki Zoo and Herpetological Society of Finland), N. Sivan (HUJ), S. Stefanni (University of Bristol), K. Talke (Bristol/Helsinki), R. Tinsley (University of Bristol), E. Wenman (Zoological Society of London), Y. L. Werner (HUJ), A. Wynn (USNM), Y. Yom-Tov (TAU), G. Zug (USNM), and all lab technicians in the above institutions.

#### LITERATURE CITED

- ANONYMOUS. 1964. International Code of Zoological Nomenclature adopted by the XV International Congress of Zoology, London, July 1958. ITZN, London.
- BAHA EL DIN, M. 1999. Egyptian tortoise conservation project report. Tortoise Trust News. Winter 1998:15.
- BAHA EL DIN, S. M. 1992. Notes on the herpetology of North Sinai. Brit. Herpetol. Soc. Bull. 41:9–11.
- . 1994. Status of the Egyptian Tortoise *Testudo kleinmanni* in Egypt. A Report on the Results of a Recent Assessment of the Conservation Status of the Species in Egypt. Turtle Recovery Programme and Wildlife Conservation Society, privately published.
- BOETTGER, O. 1893. Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. I. Teil (Rhynchocephalen, Schildkröten, Krokodile, Eidechsen, Chamäleons). Gebrüder Knauer, Frankfurt a. M., Germany.
- BRINGSØE, H., AND J. R. BUSKIRK. 1998. Distribution of *Testudo kleinmanni* Lortet, 1883 and *Testudo graeca* Linnaeus, 1758 in the Negev desert, southern Israel (Reptilia: Testudines: Testudinidae). Faun. Abh. Mus. Tierk. Dresden 21, Suppl. 4:23–30.
- BUSKIRK, J. R. 1985. The endangered Egyptian tortoise *Testudo kleinmanni*: status in Egypt and Israel.

- In S. McKeown, F. Caporaso and K. H. Peterson (eds.), Ninth Herpetological Symposium on Captive Propagation and Husbandry, p. 35–52. Univ. of San Diego, San Diego, CA.
- CALABRESI, E. 1924. Missione zoologica del Dr. E. Festa in Cirenaica. *Boll. Mus. Zool. Anat. Comp. R. Univ. Torino* 38:7–8.
- CANN, J., AND J. M. LEGLER. 1994. The Mary River tortoise: A new genus and species of short-necked chelid from Queensland, Australia (Testudines: Pleurodira). *Chelon. Conserv. Biol.* 1:81–96.
- ERNST, C. H., AND J. H. LOVICH. 1986. Morphometry in the chelid turtle, *Platemys platycephala*. *Herpetol. J.* 1:66–70.
- ERNST, C. H., J. C. WILGENBUSCH, T. P. BOUCHER, AND S. W. SEKSCIENSKI. 1998. Growth, allometry and sexual dimorphism in the Florida box turtle, *Terrapene carolina bauri*. *Herpetol. J.* 8:72–78.
- FLOWER, S. S. 1933. Notes on the recent reptiles and amphibians of Egypt, with a list of the species recorded from that Kingdom. *Proc. Zool. Soc. Lond.* 1933:734–851.
- FRITZ, U., AND J. R. BUSKIRK. 1997. The occurrence of *Testudo kleinmanni* in Tripolitania. *Časopis Národního muzea Řada přírodovědná* 167:143–144.
- FROST, D. R., AND D. M. HILLIS. 1990. Species in concept and practice: herpetological applications. *Herpetologica* 46:87–104.
- GEFFEN, E. 1990. The threatened Egyptian tortoise (*Testudo kleinmanni*): Proposal for a reserve at Holot Agur (western Negev, Israel). *Brit. Herpetol. Soc. Bull.* 32:6–8.
- GEFFEN, E., AND H. MENDELSSOHN. 1988. Home range use and seasonal movements of the Egyptian tortoise (*Testudo kleinmanni*) in the northwestern Negev, Israel. *Herpetologica* 44:254–359.
- . 1989. Activity patterns and thermoregulatory behaviour of the Egyptian tortoise *Testudo kleinmanni* in Israel. *J. Herpetol.* 23:404–409.
- . 1991. Preliminary study on the breeding pattern of the Egyptian tortoise, *Testudo kleinmanni*, in Israel. *Herpetol. J.* 1:574–577.
- GMIRA, S. 1995. Étude des Chéloniens Fossiles du Maroc. CNRS Éditions, Paris.
- GRISMER, L. L. 1999. An evolutionary classification of reptiles on islands in the Gulf of California, México. *Herpetologica* 55:446–469.
- GÜNTHER, A. 1869. Report on two collections of Indian reptiles. *Proc. Zool. Soc. Lond.* 1869:500–507.
- HIGHFIELD, A. C. 1997. 250 Egyptian tortoises in Cairo emergency. *Tortoise Trust News*. Winter 1996: 8–9.
- IYERSON, J. B. 1992. A Revised Checklist with Distribution Maps of the Turtles of the World. Privately printed. Richmond, Indiana.
- IYERSON, J. B., AND T. E. GRAHAM. 1990. Geographic variation in the redbelly turtle, *Pseudemys rubriventris* (Reptilia: Testudines). *Ann. Carnegie Mus.* 59: 1–13.
- IYERSON, J. B., AND W. P. MCCORD. 1989. The taxonomic allocation of *Emys nigricans* Gray, *Emys muticus* Cantor, and *Geoclemys kwangtungensis* Pope. *Amphib.-Reptilia* 10:23–33.
- . 1994. Variation in East Asian Turtles of the Genus *Mauromys* (Bataguridae; Testudines). *J. Herpetol.* 28:178–187.
- . 1997. A new species of *Cyclemys* (Testudines: Bataguridae) from Southeast Asia. *Proc. Biol. Soc. Wash.* 110:629–639.
- JACOBSON, E. R., M. B. BROWN, I. M. SCHUMACHER, B. R. COLLINS, R. K. HARRIS, AND P. A. KLEIN. 1995. Mycoplasmosis and the desert tortoise (*Gopherus agassizii*) in Las Vegas Valley, Nevada. *Chel. Conserv. Biol.* 1:279–284.
- LEVITON, A. E., R. H. GIBBS JR., E. HEAL, AND C. E. DAWSON. 1985. Standards in herpetology and ichthyology. Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. *Copeia* 1985:802–832.
- LORTET, L. 1883. Études zoologiques sur la faune du Lac de Tibériade suivies d'un aperçu sur la faune des lacs d'Antioche et de Homs. I. Poissons et reptiles du Lac de Tibériade et de quelques autres parties de la Syrie. *Arch. Mus. Hist. Natur. Lyon* 3:13–190.
- . 1887. Observations sur les tortues terrestres et paludines du bassin de la Méditerranée. *Arch. Mus. Hist. Natur. Lyon* 4:1–26.
- LOVERIDGE, A., AND E. E. WILLIAMS. 1957. Revision of the African tortoises and turtles of the suborder Cryptodira. *Bull. Mus. Comp. Zool. Harvard* 115: 160–557 + 18 pl.
- MERTENS, R. 1967. Die herpetologische Sektion des Natur-Museums und Forschungs-Institutes Senckenberg in Frankfurt a.M. nebst einem Verzeichnis ihrer Typen. *Senck. biol.* 48, Sonderheft A:1–106.
- MERTENS, R., AND H. WERMUTH. 1955. Die rezenten Schildkröten, Krokodile und Brückenechsen. Eine kritische Liste der heute lebenden Arten und Rassen. *Zool. Jb. (Syst.)* 83:323–440.
- MORAVEC, J., S. BAHA EL DIN, H. SELIGMANN, N. SIVAN, AND Y. WERNER. 1999. Systematics and distribution of the *Acanthodactylus pardalis* group (Lacertidae) in Egypt and Israel. *Zool. Middle East* 17: 21–50.
- PERÄLÄ, J. 2000. The Egyptian tortoise, *Testudo kleinmanni* Lortet, 1883, on the verge of extinction. *Herpetomania* 8(4):35–40 (in Finnish with English abstract).
- SCHLEICH, H. H. 1989. Merkmalsausbildungen an Landschildkröten in Nord-Ost Lybien (Testudines: Testudinidae). *Herpetozoa* 1:97–108.
- SCHLEICH, H. H., W. KÄSTLE, AND K. KABISCH. 1996. Amphibians and Reptiles of North Africa. Koeltz Scientific Books, Koenigstein, Germany.
- SEIDEL, M. E., J. M. STUART, AND W. G. DEGENHARDT. 1999. Variation and species status of slider turtles (Emydidae: *Trachemys*) in the southwestern United States and adjacent Mexico. *Herpetologica* 55:470–487.
- SIMPSON, G. G. 1961. Principles of Animal Taxonomy. Columbia Univ. Press, New York.
- TRISTRAM, H. B. 1884. The Survey of Western Palestine. Committee of the Palestine Exploration Fund, London.
- WERNER, Y. L. 1982. Georg Haas, 1905–1981. *Copeia* 1982:491–493.
- . 1987. Ecological zoogeography of the Sahara-Arabian, Saharan and Arabian reptiles in the sand deserts of southern Israel. In F. Krupp, W. Schneider, and R. Kintzelbach (eds.), Proceedings of the Symposium on the Fauna and Zoogeography of



- the Middle East, Mainz 1985, pp. 272–295. Beihefte zum TAVO A 28(1987), Mainz, Germany.
- . 1988. Herpetofaunal survey of Israel (1950–85), with comments on Sinai and Jordan and on zoogeographical heterogeneity. In Y. Yom-tov, and E. Tchernov (eds.), *The Zoogeography of Israel. The Distribution and Abundance at a Zoogeographical Crossroad*. Monographiae Biologicae Vol. 62, pp. 355–387. Dr. W. Junk Publishers, Dordrecht, The Netherlands.
- WILEY, E. O. 1978. The evolutionary species concept reconsidered. *Syst. Zool.* 29:76–80.

Accepted: 23 January 2001.

#### APPENDIX 1

##### SPECIMENS EXAMINED

*Testudo kleinmanni*. Adult males used in morphometric analyses.—LIBYA: Gheminez, Cyrenaica, USNM 139094; Cyrenaica, Derna, MTKD D 31598; NE Libya (Cyrenaica), PCHP 4319, 4321, 4323, 4340-1, 4438, 4441-3, 4445-6, 4484; Libya, MTKD D 26016, 28287, 29119. EGYPT: Alexandria, BMNH 1920.1.20.774, MG 42006132 (paralectotype), MG 42006138 (paralectotype); vicinity of Alexandria, MTKD D 32798; Mediterranean coast, MTKD D 31803, 32831; Salum, Western Desert Governate, MCZ 54045; Egypt, BMNH 1933.9.3.9-10, MNHN 7838 (ex 1879-774), MTKD D 32742, PCHP 3844, 3991, 4017, TAU 14198.

Adult females used in morphometric analyses.—LIBYA: NE Libya (Cyrenaica), MTKD D 28286, PCHP 4316-7, 4322, 4333, 4335-6, 4337-8, 4435-7, 4439, 4440, 4447-54; Gheminez, Cyrenaica, USNM 139092; ca 30-50km W Tripoli, close to coast, MTKD D 30855; W Libya (Tripolitania), MTKD D 32434-5, 34854; Libya, MTKD D 26017, 32832, PCHP 4686. EGYPT: Alexandria, BMNH 88.4.24.2, 1920.1.20.773, MG 42000413 (paralectotype), MG 42000414, designated lectotype of *Testudo kleinmanni* (this article), MG 42006125 (ex MG 110) (paralectotype); neighborhood of Alexandria, BMNH 1897.11.13.2-3 (only 1 specimen: male apparently lost); Mersa Matruh, NW Egypt, BMNH 1933.9.3.16; Agniba near Mersa Matruh, BMNH 1977.1218; Egypt, BMNH 1933.9.3.7, MNHN 1878-602 (DA 51 A) (paralectotype), 1987-971 (DA 51 B) (paralectotype), PCHP 3843, 3954-6; "Sindh" (extralimital), BMNH 1947.3.4.35 (ex BMNH 69.8.28.5), presented by Dr. Leith 1869 (type of *Testudo leithii* Günther, 1869).

Additional specimens examined.—EGYPT: Alexandria, MCZ 5081 (paralectotype), juv., MG 42006110 (paralectotype), deformed male; Giza, USNM 55758, adult male, no gular; Egypt, MNHN 1876-416 (paralectotype), skull, MNHN 1992.197 (DD 62) (paralectotype), skull and misc. bones; MNHN 9462 (paralectotype) (juv. female).

*Testudo wernerii*. Adult males used in morphometric analyses.—EGYPT: Suez Canal, Kantarah-il-Khastreh, BMNH 1900.2.8.11 (paratype); N Sinai, BMNH 1933.9.3.12 (paratype); Egypt, BMNH 1933.9.3.6. ISRAEL: Negev: ca. 14 km S Be'er Sheva, HUI 16047 (paratype), HUI 16048 (paratype); Be'er Sheva, TAU 14120 (paratype); near Ashalim, Negev desert, JP 97.3.36 (examined in the field, data deposited in BMNH); 3 km N of Revivim, TAU 5636 (paratype); Gevulot, TAU 7468 (paratype), TAU 11941 (paratype); 1 km E of Rehovot, TAU 8363 (paratype); Agur Sands, TAU 12022 (paratype); Holot Agur, TAU 12769 (paratype); Holot Nahagur, TAU 12712 (paratype); Nahal Secher, TAU 14126 (paratype).

Adult females used in morphometric analyses.—EGYPT: N Sinai, BMNH 1933.9.3.14 (paratype); Egypt, BMNH 1933.9.3.5. ISRAEL: sand dunes 14 km S of Be'er Sheva, HUI 949, coll. Haas, Werner et al. 12.3.1963 (first record of *T. kleinmanni* in Israel; holotype of *T. wernerii* sp. nov.); 14 km S Be'er Sheva, HUI 989 (paratype); 16 km S of Be'er Sheva, TAU 11898 (paratype); near Ashalim, Negev desert, JP 97.3.37 (examined in the field, data deposited in BMNH); Tel-Yerucham, TAU 6295 (paratype); Agur, TAU 13397 (paratype).

Additional specimens examined.—EGYPT: Sinai: El Bardawil, west to Nahal Yam settlement, HUI 971 (crushed ad. male); North Sinai, HUI 16142 (crushed ad.). ISRAEL: Bir Asluj sands, Boaz Shacham, 4 July 1999, HUI (uncatalogued hatchling); Be'er Chagil, TAU 12017 (ad. female, plastron deformed); Kibutz Gevulot, TAU 13150 (juv. male); Agur Sands, HUI 22820, (juv. female), TAU 13889 (juv. male); Tel-Aviv, MNHN 1984-500 (deformed juv.), extralimital: probably from TAU's Expo Zoo, don. Mendelssohn; Israel (originally wild caught captives from TAU's Expo Zoo), TAU 11103 (juv. male), 11129 (juv.), 12141 (adult male), 12216 (adult male), 13139 (adult male), 13952-3 (juv. males), 14121 (ad. female), 14122-3 (ad. males), 9359 (ad. male), 9360-1 (ad. females).

Putative Hybrids.—West Libya, MTKD D 34854; Libya: Gheminez, Cyrenaica USNM 139093; Gevulot, Israel, TAU 12026; S of Gevulot, Israel, TAU 5635.